

EYE CHART

Presbyopia Patient Directions

View the eye chart without reading glasses. Take product as directed by your eye doctor. View the eye chart again.



Talk to your eye doctor for more information.

Font size¹

Vision Test

16 pt	Presbyopia is a natural part of aging.
14 pt	You're likely not the only person reading
12 pt	an eye chart like this right now. The first
9.5 pt	symptoms most people notice are difficulty
7.5 pt	reading fine print, especially in low light conditions.
6 pt	Difficulty seeing and focusing on objects that are close to you
4.5 pt	is another common symptom. If you need brighter lighting or have to hold
3.5 pt	materials at an arm's distance to see them clearly, you could be experiencing symptoms.
3 pt	<small>If you're squinting a lot when reading up close you could be experiencing a common symptom.</small>

USE

VUITY[™] (pilocarpine hydrochloride ophthalmic solution) 1.25% is a prescription eye drop used to treat age-related blurry near vision (presbyopia) in adults.

IMPORTANT SAFETY INFORMATION

- Do not use VUITY if you are allergic to any of the ingredients.
- Use caution when driving at night or performing hazardous activities in poor lighting.
- Temporary problems when changing focus between near and distant objects may occur. Do not drive or use machinery if vision is not clear.
- Seek immediate medical care if you experience any sudden vision loss.

Please see additional Important Safety Information on the following page and the full [Prescribing Information](#).



DROP BY TO SEE MORE
at Vuity.com

IMPORTANT SAFETY INFORMATION (CONTINUED)

- If you wear contact lenses, they should be removed prior to VUITY use. Wait 10 minutes after dosing before reinserting contact lenses.
- Do not touch the dropper tip to any surface as this may contaminate the contents.
- If more than one topical eye medication is being used, the medicines must be administered at least 5 minutes apart.
- The most common side effects are headache and eye redness. These are not all the possible side effects of VUITY.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

Please see Use and additional Important Safety Information on the previous page and the full [Prescribing Information](#).

Reference: 1. Sanders DR, Sanders ML. J Refract Surg. 2007;23:747-751.



VUITY™ and its design are registered trademarks of Allergan, Inc., an AbbVie company.
© 2021 AbbVie. All rights reserved. US-VUI-210174 December 2021